

Aberystwyth University

Microbially enhanced carbon capture and storage by mineral-trapping and solubility-trapping

Mitchell, Andrew C.; Dideriksen, Knud; Spangler, Lee H.; Cunningham, Alfred; Gerlach, Robin

Published in:
Environmental Science & Technology

DOI:
[10.1021/es903270w](https://doi.org/10.1021/es903270w)

Publication date:
2010

Citation for published version (APA):
Mitchell, A. C., Dideriksen, K., Spangler, L. H., Cunningham, A., & Gerlach, R. (2010). Microbially enhanced carbon capture and storage by mineral-trapping and solubility-trapping. *Environmental Science & Technology*, 44(13), 5270-5276. <https://doi.org/10.1021/es903270w>

General rights

Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400
email: is@aber.ac.uk

Microbially Enhanced Carbon Capture and Storage by Mineral-Trapping and Solubility-Trapping

ANDREW C. MITCHELL,^{*,†,§}
 KNUD DIDERIKSEN,[§] LEE H. SPANGLER,[†]
 ALFRED B. CUNNINGHAM,[†] AND
 ROBIN GERLACH[†]

Center for Biofilm Engineering, Montana State University,
 Bozeman, Montana, 59717, Department of Chemistry and
 Biochemistry, Montana State University, Bozeman, Montana,
 59717, and NanoGeoScience Group, Nano-Science Center,
 Department of Chemistry, University of Copenhagen,
 DK-2100 Copenhagen Ø, Denmark

Received November 4, 2009. Revised manuscript received
 March 24, 2010. Accepted April 6, 2010.

The potential of microorganisms for enhancing carbon capture and storage (CCS) via mineral-trapping (where dissolved CO₂ is precipitated in carbonate minerals) and solubility trapping (as dissolved carbonate species in solution) was investigated. The bacterial hydrolysis of urea (ureolysis) was investigated in microcosms including synthetic brine (SB) mimicking a prospective deep subsurface CCS site with variable headspace pressures [p(CO₂)] of ¹³C–CO₂. Dissolved Ca²⁺ in the SB was completely precipitated as calcite during microbially induced hydrolysis of 5–20 g L⁻¹ urea. The incorporation of carbonate ions from ¹³C–CO₂ (¹³C–CO₃²⁻) into calcite increased with increasing p(¹³CO₂) and increasing urea concentrations: from 8.3% of total carbon in CaCO₃ at 1 g L⁻¹ to 31% at 5 g L⁻¹, and 37% at 20 g L⁻¹. This demonstrated that ureolysis was effective at precipitating initially gaseous [CO₂(g)] originating from the headspace over the brine. Modeling the change in brine chemistry and carbonate precipitation after equilibration with the initial p(CO₂) demonstrated that no net precipitation of CO₂(g) via mineral-trapping occurred, since urea hydrolysis results in the production of dissolved inorganic carbon. However, the pH increase induced by bacterial ureolysis generated a net flux of CO₂(g) into the brine. This reduced the headspace concentration of CO₂ by up to 32 mM per 100 mM urea hydrolyzed because the capacity of the brine for carbonate ions was increased, thus enhancing the solubility-trapping capacity of the brine. Together with the previously demonstrated permeability reduction of rock cores at high pressure by microbial biofilms and resilience of biofilms to supercritical CO₂, this suggests that engineered biomineralizing biofilms may enhance CCS via solubility-trapping, mineral formation, and CO₂(g) leakage reduction.

Introduction

Geologic sequestration of carbon dioxide, also known as carbon capture and storage (CCS), is one strategy to reduce the emission of greenhouse gases generated through the combustion of fossil fuels. Geologic sequestration of CO₂ involves the injection of supercritical CO₂ (SC–CO₂; critical point = 31.1 °C and 73 atm) into underground formations such as oil bearing formations, deep unmineable coal seams, and deep saline aquifers (1, 2). Methods that can trap CO₂ in a nonlabile phase would be advantageous in the development of subsurface CO₂ storage as a viable engineered mechanism to reduce the net emission of CO₂ from fossil fuel combustion (2–5). This includes so-called mineral-trapping, where carbonate ions from the injected CO₂ can be incorporated into a solid carbonate mineral, and solubility-trapping, where aqueous inorganic carbon species are contained in the formation water itself (6). SC–CO₂ injection into deep underground formations can result in elevated pressure in the region surrounding the point of injection. As a result, an upward hydrodynamic pressure gradient may develop across the trapping cap-rock, leading to the upward leakage of CO₂, due to the primary permeability of the cap-rock, through fractures, or near injection wells (2). Therefore, methods that can simultaneously reduce such leakage and facilitate trapping CO₂ in nonlabile forms would be advantageous. Calcium is the most abundant cation in many surface- and ground-waters (7) and therefore mineralization in CaCO₃, a common rock forming mineral, may provide an effective means of immobilization during mineral-trapping. Equally, amendments to brine chemistry may enhance the capacity of the brine for CO₂(g) and dissolved carbonate ions, thus increasing the solubility-trapping capabilities.

Laboratory and field studies into SC–CO₂–fluid–rock interactions are limited, but general processes have been investigated for injection into brine formations (8). Injection of SC–CO₂ into brine aquifers initially reduces the pH of the brine by 1.5–4 pH units due to the dissolution of CO₂ and disassociation of H₂CO₃ (eqs 1–4) (7) depending on the brine chemistry, formation lithology, and temperature (9). However, carbonate alkalinity is also produced by the reaction of H₂CO₃ with reservoir minerals (8–10), and computational and laboratory experiments suggest that overall, pH actually increases by around 0.5 to 1 units relative to the preinjection brine (9).

Mineralization of injected CO₂ into CaCO₃ will be possible if the equilibrium of the reaction $\text{Ca}^{2+} + \text{CO}_3^{2-} \rightleftharpoons \text{CaCO}_3$ can be moved to the right and the saturation state of CaCO₃ is exceeded. This may be achievable if sufficient dissolved Ca is present, if pH and alkalinity can be increased, and if appropriate nucleation substrates are present (7). Microbiologically enhanced CaCO₃ precipitation offers a potential mechanism to achieve this in situ. Microbes have been shown to enhance CaCO₃ precipitation via cation adsorption to negatively charged functional groups on microbe surfaces and by metabolically driven changes in the solution chemistry, which increase mineral saturation and induce nucleation (11). An increase in pH, CO₃²⁻, and HCO₃⁻ can be caused for instance by bacterial ureolysis (the enzymatic hydrolysis of urea) (12), sulfate, iron, or nitrate reduction (13, 14), or alkalization during photosynthesis (15). In subsurface environments, CaCO₃ precipitation induced by the ureolysis of injected urea has been investigated as a potential method to reduce bedrock porosity for enhanced oil recovery (16) and for coprecipitating radionuclides from groundwater in the vadose zone of contaminated aquifers (12, 17–19).

* Corresponding author e-mail: andrew.mitchell@erc.montana.edu.


[†] Center for Biofilm Engineering, Montana State University.

[§] University of Copenhagen.


[‡] Department of Chemistry and Biochemistry, Montana State University.

Ureolysis results in the production of ammonium (NH_4^+), dissolved inorganic carbon (DIC), and an increase in pH via a net production of OH^- ions (eqs 5–9), which favors CaCO_3 precipitation (eqs 10 and 11).


$\text{CO}_2(\text{g})$ Dissolution and Disassociation.



Ureolysis.



Overall Ureolysis and CaCO_3 Precipitation.



The ability of ureolysis to operate in dark subsurface environments, and the high rates of ureolysis induced CaCO_3 precipitation relative to sulfate reduction induced CaCO_3 precipitation (13, 18), provides a viable mechanism to induce subsurface CaCO_3 precipitation and CO_2 mineralization. Additionally, the associated reduction in porosity in subsurface environments would reduce the potential of CO_2 leakage after injection has ceased. The change in brine chemistry during ureolysis may also enhance the solubility-trapping capacity of the brine through pH changes and thus carbonate ion speciation (7). Additionally, microbial biofilms, which are assemblages of microorganism attached to a surface, have shown remarkable resilience to SC-CO_2 . This suggests microorganisms in a biofilm form may offer physical, chemical, and biological mechanism to enhance CCS under high pressure during and after injection of SC-CO_2 (4, 5) (Figure 1).

Here the effect of bacterially induced ureolysis on CaCO_3 precipitation was investigated for a range of fixed initial CO_2 headspace pressures and urea concentrations in synthetic brine mimicking the Powder River Basin in Wyoming, a prospective CO_2 injection site in the western U.S. (20). Lower than reservoir pressure conditions are considered in this study in order to determine the general relationship between ureolysis, mineral formation, and carbonate sources, as well as to ensure the applicability of modeling results, for which accurate thermodynamic data exists.

Experimental Methods

Experimental Setup. Batch CO_2 precipitation microcosms were prepared in sterile 35 mL serum bottles in a laminar flow hood. Each microcosm included 20 mL of a synthetic brine (SB) with approximately 5 g L^{-1} total dissolved solids (TDS), mimicking the ionic ratios present in the proposed Powder River Basin CO_2 injection site (20). Specific inorganic constituents of the SB are shown in the Supporting Information (SI) Table 1. Three SBs were produced with different




FIGURE 1. Schematic representation of microbially enhanced carbon capture and storage.

urea [$(\text{NH}_2)_2\text{CO}$] concentrations of 1, 5, or 20 g L^{-1} , to simulate the addition of varying concentrations of urea to the aquifer. Test and control SB microcosms were prepared. Test SB microcosms included the ureolytic organism *Sporosarcina pasteurii* (ATCC 11859, gram-positive, spore-forming, urease positive), known to induce CaCO_3 precipitation (12, 17, 19). Spore forming bacilli have been shown to be resistant to high pressures and the exposure to SC-CO_2 (4, 5, 21). Biotic control experiments included the nonureolytic gram-positive bacterium *Bacillus subtilis* strain 186 (ATCC 23857). Abiotic control experiments included just the SB with no bacterial amendment. The headspace (15 cm^3) of each vial was then purged with 0.2 μm filter sterilized 99% $\delta^{13}\text{C}$ - CO_2 (Cambridge Isotope Laboratories, MA) for 1 min, and final headspace $p(\text{CO}_2)$ of 1, 1.35, and 1.7 atm were attained by measurement with a headspace pressure gauge. A set of microcosms was also retained with an atmospheric pressure air headspace [390 ppm $\text{CO}_2(\text{g})$ at 1 atm = 0.00039 atm $p(\text{CO}_2)$]. The microcosms were incubated at 22 $^\circ\text{C}$ on a shaker at 100 rpm for 8 days. While headspace pressures are far lower than expected during deep subsurface SC-CO_2 injection (>73 atm) (8), these microcosms allow investigation of the general relationship of $p(\text{CO}_2)$ and ureolysis on carbonate speciation. Additionally, even in deep subsurface brines the $p(\text{CO}_2)$ will likely vary significantly due to the advective and diffusive transport of CO_2 from the injection point over time (22). The SB was characterized for pH, dissolved Ca, and NH_4^+ immediately after inoculation and headspace adjustment [time (t) = 0, t_0], and again after 8 days (t_8) (See SI). At the termination of the experiments any CaCO_3 precipitates formed in the microcosms were characterized by X-ray diffraction system (XRD) and analyzed for $\delta^{13}\text{C}$ (SI, eq 1).

Speciation and Saturation Modeling. PHREEQC (23) was used to model inorganic carbon speciation and carbonate mineral saturation in microcosm experiments simulating CO_2 injection into the Powder River Basin brine with $p(\text{CO}_2)$ of 0.00039, 1, and 1.7 atm. All calculations were carried out at 25 $^\circ\text{C}$ and used the MINTEQ database (24), to which the thermodynamic constants for urea was added (25) (see SI). PHREEQC calculates the saturation index (S) of the reacting solution with respect to calcite according to

$$S_{\text{calcite}} = \log \frac{\{\text{Ca}^{2+}\}\{\text{CO}_3^{2-}\}}{K_{\text{SO}}} \quad (13)$$

where $\{\text{Ca}^{2+}\}$ and $\{\text{CO}_3^{2-}\}$ are the activities of dissolved Ca^{2+} and CO_3^{2-} , and K_{SO} is the equilibrium calcite solubility product (7).

Results and Discussion

Synthetic Brine Chemistry. At t_0 , the biotic test (*S. pasteurii*), biotic control (*B. subtilis*), and abiotic control experiments under different $p(\text{CO}_2)$ and initial urea concentrations exhibited starting chemistries consistent with the SB (SI Table 1), with dissolved Ca concentrations of ~ 17 mM (Figure 2; SI, Table 2). NH_4^+ concentrations were ~ 0.4 mM, likely due to some limited abiotic urea hydrolysis. The initial pH of the SB with $p(\text{CO}_2)$ of 0.00039 atm was ~ 6 , with lower pH (between 4 and 5) exhibited in the SB with higher $p(\text{CO}_2)$ (1 to 1.7 atm), consistent with a higher concentration of protons liberated from the disassociation of carbonic acid at higher $p(\text{CO}_2)$ (Figure 2; SI, Table 2) (7). The SB at t_0 was undersaturated with respect to calcite at all $p(\text{CO}_2)$. The predicted calcite saturation index (S_{calcite}) decreased from approximately -1 to -4 with $p(\text{CO}_2)$ increasing from 0.00039 to 1.7 atm (Figure 3).

After 8 days (t_8), biotic control and abiotic control experiments under different $p(\text{CO}_2)$ and initial urea concentrations demonstrated very little, if any, NH_4^+ production associated with some limited abiotic urea hydrolysis, but the pH remained largely unchanged (Supporting Information, Table 2). Dissolved Ca concentrations remained unchanged relative to t_0 , indicating no CaCO_3 precipitation in biotic and abiotic control experiments, consistent with the unchanged and undersaturated solution from t_0 . This would be expected with the biotic control inoculated with the non ureolytic organism, *B. subtilis*, or with the abiotic control without bacterial amendment, as has been observed previously (12, 17, 18, 26).

Conversely, in the biotic test experiments, NH_4^+ concentrations increased, consistent with the hydrolysis of urea by *S. pasteurii* (12, 17, 27, 28). All available urea was hydrolyzed in the 1 g L⁻¹, 5 g L⁻¹, and 20 g L⁻¹ experiments, under different $p(\text{CO}_2)$, which is consistent with previous studies using up to 20 g L⁻¹ (12, 16, 26, 28). The pH also increased in the *S. pasteurii* inoculated experiments, with relative increases in pH during the experiments proportional to the amount of urea hydrolyzed, and thus proportional to the initial urea concentrations (Figure 2).

Calcium Carbonate Precipitation in Synthetic Brine at Different $p(\text{CO}_2)$. Dissolved calcium concentrations decreased by between 97 and 100% in the biotic experiments with $p(\text{CO}_2)$ of 1, 1.3, and 1.7 atm and with 5 or 20 g L⁻¹ urea (Figure 2; SI, Table 2). Dissolved Ca reductions were slightly lower, at between 87 and 98% in the biotic experiments with $p(\text{CO}_2)$ of 0.00039 atm and 5 or 20 g L⁻¹ urea. This was consistent with the precipitation of up to 17 mM CaCO_3 (Figure 3). Biotic test experiments with 1 g L⁻¹ urea exhibited only a slight decrease in dissolved Ca at elevated $p(\text{CO}_2)$ (Figure 2; SI, Table 2) but significant reduction in dissolved Ca at 0.00039 atm. This precipitation of CaCO_3 in the 0.00039 atm treatments is explained by the initially higher saturation index compared to the treatments with higher initial $p(\text{CO}_2)$ (Figure 3).

XRD analysis confirmed the CaCO_3 polymorph precipitated was calcite, although small amounts of vaterite and aragonite were morphologically observable by scanning electron microscopy (SEM) (data not shown).

Inorganic Carbon Speciation in CaCO_3 and Artificial Brine. $\delta^{13}\text{C}$ - CaCO_3 was used to discriminate the origin and proportion of carbonate ions in precipitated calcite under




FIGURE 2. Synthetic brine ammonium (NH_4^+) concentration, pH, and Ca^{2+} concentration in CO_2 biomineralization experiments inoculated with *Sporosarcina pasteurii*, at varying $\text{CO}_2(\text{g})$ headspace pressures and urea concentrations at time t_0 (0 days) and t_8 (8 days). Error bars smaller than symbol if not visible.

different $p(\text{CO}_2)$ and urea concentrations (SI, Table 3). Urea and ^{13}C - $\text{CO}_2(\text{g})$ provide two potential carbonate ion sources in the microcosm experiments simulating CO_2 injection into the Powder River Basin brine aquifer. The ^{13}C - CO_2 had a $\delta^{13}\text{C}$ of $+89000$ ‰ (= 99% ^{13}C - CO_2 , see SI, eq 2 for details), and the urea used exhibited a $\delta^{13}\text{C}$ of -40.6 ‰. The SB also




FIGURE 3. Synthetic brine CaCO₃ (calcite) saturation index (S_{CaCO_3}) at time t_0 and mass of CaCO₃ precipitated (mM) at time t_0 and t_8 (8 days) at varying CO₂(g) headspace pressures and urea concentrations in CO₂ biomineralization experiments inoculated with *Sporosarcina pasteurii*.

contained 0.25 mM MgCO₃ with a $\delta^{13}C$ of -8.0 ‰. The $\delta^{13}C$ -CaCO₃ of calcite precipitated from the 1, 1.3, and 1.7 atm p(CO₂) microcosms ranged from +4.0 to +57877 ‰, indicating the precipitated calcite was heavily enriched with carbon originally from the ¹³C-CO₂(g) over the brine (SI, Table 3).

Natural isotopic equilibrium fractionation can be expected during CO₂(g) dissolution and disassociation, between CO₂(g) and dissolved bicarbonate, but this will only account for an increase of +7.9 to +8.4 ‰ (29). Calcite precipitated with 0.00039 atm p(CO₂) exhibited $\delta^{13}C$ -CaCO₃ of +1.1, reflecting the natural abundance of ¹³C on Earth, and consistent with most natural calcites precipitated at ambient p(CO₂) (29). These data demonstrate that the ¹³C as a % of total C in CaCO₃ increased with increasing p(CO₂) for each of the urea concentrations (Figure 4a). Additionally, higher urea concentrations increased the proportion of ¹³C incorporated into CaCO₃ at a given p(CO₂), from an average of 8.3% at 1 g L⁻¹ to 31% at 5 g L⁻¹ and 37% at 20 g L⁻¹ (Figure 4a). A maximum of 65% of carbon from the original atmosphere of the microcosm was precipitated in calcite. It can be seen from the stoichiometry of the ureolysis reaction (eqs 10 and 11), that no net precipitation of CO₂(g) in CaCO₃ occurred during urea hydrolysis-induced mineral-trapping. However, the increase in $\delta^{13}C$ -CaCO₃ with increasing p(CO₂) and at higher urea concentrations with equivalent p(CO₂) demonstrates that (i) ureolysis enhances the proportion of carbonate ions precipitated in CaCO₃ that are from the originally gaseous CO₂ and (ii) there is a higher proportion of carbonate ions from the originally gaseous ¹³CO₂ relative to carbonate ions derived from urea dissolved in the brine. Previous studies suggest that the C isotopes of dissolved carbonate and CO₂(g) equilibrate within 12 to 24 h (30, 31). This shows that extensive transport of ¹³C from the gas phase to the aqueous phase would occur without a net flux of carbon from the gas phase.




FIGURE 4. (a) Measured and (b) modeled ¹³C as a percentage of total C in precipitated CaCO₃ in CO₂ biomineralization experiments inoculated with *Sporosarcina pasteurii* by t_8 (8 days). Dashed lines represent models which include 50 mg L⁻¹ bacterial organic carbon by t_8 (8 days).

Therefore, the CO₂(g) concentration in the gas phase was unlikely to be lowered by up to 65%, as might be suggested from the isotope data. Thermodynamic modeling on PHREEQC (23) allowed testing of the validity of the experimental results, in order to examine the speciation of carbon from CO₂(g) to aqueous species during ureolysis induced carbonate precipitation in brine.

The modeled pH and mass of calcite precipitated by t_8 (SI, Table 3) is largely in agreement with the experimental values. The actual amount of Ca scavenged in experiments with 1 g L⁻¹ urea is slightly larger than in the calculations. Possibly, this may reflect Ca adsorption to bacterial surfaces in the experiments, as observed previously (32), an effect that has not been included in the model. From PHREEQC calculations, the ¹³C content of the precipitated calcite was predicted based on the assumption that the equilibration of C isotopes in solution and gas phase occurred at rates much faster than the calcite precipitation. Previous studies demonstrate precipitation occurs over 2–3 days (12), versus equilibration of C isotopes in solution and gas phase which occurs within 12 to 24 h (30, 31), so the assumption is valid. For each step of urea hydrolysis, the sequence of the calculation was (i) addition of 1 mM carbonate with $\delta^{13}C = 0$ ‰ to the solution to reflect urea hydrolysis; (ii) equilibration of the C isotopes of the solution and the gas phase [initial $\delta^{13}CO_2 = 89\ 000$ ‰] based on mass balance; (iii) when calcite precipitated from supersaturated solutions, its C isotope composition reflected that of the dissolved carbonate. At the end of urea hydrolysis, when 100 mM of urea had been hydrolyzed, the final C isotope composition of the calcite was calculated from the mass and isotope composition of the solid precipitated at each step.

The modeled C isotope compositions are more positive than those determined experimentally, especially for the experiments with 1 g L⁻¹ urea (Figure 4b; SI, Table 3). The difference between measured and predicted values correlates with the amount of calcite precipitated in the experiments,

where experiments precipitating only a small mass of calcite display very light isotope compositions relative to that measured. A likely explanation for this is that all isotope analyses are influenced by carbon contamination from the bacterial biomass, that accumulated on the crystal surfaces during filtration, or within the crystals, as demonstrated previously (12). Such organic carbon is isotopically lighter with respect to carbon, therefore exhibiting a lower $\delta^{13}\text{C}$. PHREEQC modeling suggested the discrepancy between measured and modeled $\delta^{13}\text{C}$ values could be accounted for by the mass of bacterial carbon in the experiments (50 mg L^{-1}) (Figure 4b; SI, Table 3). However, it is unclear if such an extensive digestion of organic material by the standard phosphoric acid digestion used during C isotope analysis of carbonate is possible (33).

In addition, in all calculations outlined here, it was assumed that dissolved carbonate and $\text{CO}_2(\text{g})$ are at isotopic equilibrium. It is possible that this is not the case and that C isotopes of dissolved carbonate only equilibrated partly with the atmosphere during the initial calcite precipitation. Because the hydrolysis of urea is completed faster in the 1 g L^{-1} experiments, such an effect would influence results from these experiments the most. Indeed, the modeled values differ mostly from the measured values for the 1 g L^{-1} experiments. However, without detailed knowledge of the kinetics of C isotope exchange and urea hydrolysis, we cannot quantify the effect of partial isotope equilibration. Nevertheless, the combination of a feasible mass of organic carbon in the experiments, and partial equilibration of organic carbon and dissolved carbonate and $\text{CO}_2(\text{g})$ appear to account for the observed $\delta^{13}\text{C}$ of the CaCO_3 .

Evolution of Brine Chemistry and Carbonate Speciation As a Function of Ureolysis Progress. The capacity of the brine for carbonate ions and the precipitation of CaCO_3 is controlled by the complex interplay of carbonate equilibrium and pH (7). We therefore evaluated the detailed evolution of the brine chemistry and carbonate precipitation using PHREEQC (23) in order to elucidate the detailed changes in carbonate speciation and fluxes between the gas, solid, and solution phases. This was undertaken for the Powder River SB chemistry used in the microcosm experiments, including 17 mM Ca (SI, Table 1) at 0.00039 , 1.0 , and $1.7 \text{ atm } p(\text{CO}_2)$. This was also undertaken for other theoretical groundwater, based upon the chemistry of the Powder River Basin, but including Ca concentrations of 2 mM , equivalent to many common groundwater compositions (34, 35) and 125 mM Ca , equivalent to many oil field brines (36, 37) (Figure 5; SI, Figure 1).

The modeling for solutions including 17 mM Ca demonstrate that after equilibration of the groundwater solutions with the headspace $\text{CO}_2(\text{g})$, for $p(\text{CO}_2)$ above atmospheric pressure, (i) ureolysis increases pH, via the net production of hydroxide ions and consumption of protons (eq 8). (ii) Once supersaturated, CaCO_3 precipitates, and the degree of ureolysis required for the induction of precipitation is greater at higher initial $p(\text{CO}_2)$ because it takes longer for the pH to increase to the point where the solution is saturated with respect to calcite. In addition, at higher initial $p(\text{CO}_2)$, the proportion of carbonate ions precipitated in CaCO_3 decreases at a given degree of ureolysis, with a greater proportion remaining as aqueous carbonate ions $[\text{CO}_3(\text{aq})]$. (iii) Headspace $p(\text{CO}_2)$ decreases, as long as ureolysis continues and pH is increasing, as the carbonate ion capacity and aqueous carbonate ion concentration increases in response to the increasing pH. Here, at higher initial $p(\text{CO}_2)$, the mass of CO_2 dissolving into the groundwater solutions, and thus the concentration of $\text{CO}_3(\text{aq})$, increases, at a given degree of ureolysis. This reduced the headspace concentration of CO_2 by up to 32 mM per 100 mM urea hydrolyzed (Figure 5). Modeling results for calcium concentrations of 2 mM and




FIGURE 5. Modeled evolution of Powder River Basin brine chemistry and carbonate speciation as a function of ureolysis progress at variable $p(\text{CO}_2)$ in atm for a brine containing 17 mM of dissolved calcium and consistent with that used in the experimental study (SI, Table 1). Dissolved Inorganic Carbon (DIC) represents all dissolved carbonate species.

125 mM demonstrate the same general trends (SI, Figure 1). However, increased calcium concentrations lower the pH attained at a given degree of ureolysis due to the precipitation of CaCO_3 , which results in the liberation of protons during the conversion of HCO_3^- to CO_3^{2-} (eqs 4 and 11). Therefore at higher calcium concentrations, a greater mass of urea has to be hydrolyzed to attain a particular pH and DIC concentration, via the flux of $\text{CO}_2(\text{g})$ into solution as the carbonate ion capacity and aqueous carbonate ion concentration increases in response to the increasing pH.

The Potential of Microbially Enhanced CCS. The data presented here suggest the potential of microbial ureolysis to increase the rate of mineral-formation and enhance solubility-trapping of CO_2 injected into brine aquifers. This occurs primarily via the pH increase induced by ureolysis which induces carbonate precipitation and increases the carbonate capacity of the SB (7).

While CaCO_3 is precipitated, the moles of calcite precipitated were equal to or less than the moles of urea derived carbonate ions. Thus, no net precipitation of $\text{CO}_2(\text{g})$ in CaCO_3 occurred during urea hydrolysis-induced mineral-trapping. Additionally, the consumption of Ca^{2+} , and the likelihood that this will not be replenished via equilibration because of the concurrent rise in pH, may reduce the magnitude of natural carbonation in the aquifer. However, the precipitation

of CaCO_3 by ureolysis in the brine provides a number of distinct engineering advantages. First, ureolytic organisms appear to be ubiquitous in surface and subsurface soils (19, 38) and *S. pasteurii* has been shown by us to be ureolytically active at pressures and temperatures relevant to geologic carbon sequestration scenarios ($P > 89$ bar, $T > 32$ °C) (39). Thus engineering solutions could likely rely on the stimulation of native organisms to induce CaCO_3 precipitation. Second, wastewater provides a potential supply of waste urea. If wastewater urea sources can be utilized in this process, it may provide the simultaneous and advantageous degradation of urea waste, and thus a reduction in labile earth surface carbon, and the mineral- and solubility-trapping of injected CO_2 . The annual volume of wastewater was $1347 \text{ km}^3 \text{ yr}^{-1}$ in Europe, North America and Asia in 1995 (40). Wastewater commonly includes urea concentrations of 20 g L^{-1} (333 mM) (41), which is more than required to maintain calcite supersaturation in groundwater with chemistries like those of the Powder River Basin, and to precipitate out all available dissolved Ca even in concentrated oilfield brines with Ca concentrations of 125 mM or more (SI, Figure 1). Third, the precipitation of CaCO_3 in the subsurface provides a means to decrease formation porosity and reduce the potential of CO_2 leakage. For example, subsurface ureolysis induced carbonate precipitation has been investigated for permeability reduction for enhanced oil recovery and radionuclide contaminant sequestration (16).

The main potential limitation of microbially enhanced CCS is the ability of microorganisms to withstand high pressure and SC- CO_2 . Planktonic cells (free floating) show limited resistance to SC- CO_2 . Twenty-two tested vegetative species of microorganisms reported in the literature were completely deactivated at some combination of pressure and temperature in the presence of SC- CO_2 (21). However, biofilms, which are microorganism assemblages firmly attached to a surface, appear to exhibit a higher resistance to SC- CO_2 (4, 5). For example, a 19 min exposure to 35°C, 136 atm SC- CO_2 resulted in a 3 \log_{10} viability reduction of planktonic *Bacillus mojavensis* cells, but resulted in only a 1 \log_{10} reduction in viable cell numbers from biofilm cultures. It is hypothesized that the small reduction in the viability of biofilm microorganisms reflects the protective role of the biofilm extracellular polymeric substances (4, 5). The resilience of microorganisms in biofilm states, to high pressure gaseous and SC- CO_2 suggests microbially enhanced mineral-trapping and solubility-trapping of CO_2 during CCS may be effectively used. In conclusion, a microbial ureolysis based approach appears to offer potential for enhancing CCS by (i) increasing the flux of gas into the brine, and the capacity of brine for carbonate ions, and (ii) the formation of carbonate minerals, potentially reducing formation porosity and the potential of CO_2 leakage to the surface. The apparent resilience of biofilm-organisms to SC- CO_2 suggests these and other microbially mediated processes may offer the ability to enhance the capacity and rates of CO_2 trapping.

Acknowledgments

Funding was provided from the U.S. Department of Energy (DOE) Zero Emissions Research Technology Center (ZERT), Award No. DE-FC26-04NT42262 and DOE EPSCoR Award No. DE-FG02-08ER46527 as well as a European Union (EU) Marie Curie Postdoctoral Fellowship to ACM. Support for the Environmental and Biofilm Mass Spectrometry Facility through DURIP, Contract Number: W911NF0510255 and the MSU Thermal Biology Institute from the NASA Exobiology Program Project NAG5-8807 is acknowledged.

Supporting Information Available

1. Details regarding the experimental methods for the microcosm experiments, 2. Information on experimental details and calculations regarding the ^{13}C fraction in the

precipitated carbonates, 3. Information regarding the geochemical modeling. A detailed explanation on how the amount of bacterial organic carbon in filtered CaCO_3 samples was determined. Three tables and one additional figure are also included. This material is available free of charge via the Internet at <http://pubs.acs.org>.

Literature Cited

- (1) Zakkour, P.; Haines, M. Permitting issues for CO_2 capture, transport and geological storage: A review of Europe, USA, Canada and Australia. *Int. J. Greenhouse Gas Control* **2007**, *1*, 94–100.
- (2) White, C. M.; Strazisar, B. R.; Granite, E. J.; Hoffman, J. S.; Pennline, H. W. Separation and capture of CO_2 from large stationary sources and sequestration in geological formations—Coalbeds and deep saline aquifers. *J. Air Waste Manage. Assoc.* **2003**, *53* (6), 645–715.
- (3) Metz, B.; Davidson, O.; de Coninck, H.; Loos, M.; Meyer, L. (Eds.). *Carbon Dioxide Capture and Storage. Summary for Policy Makers and Technical Summary*. IPCC, Geneva, Switzerland, pp 53.
- (4) Mitchell, A. C.; Phillips, A.; Hiebert, R.; Gerlach, R.; Spangler, L.; Cunningham, A. B. Biofilm enhanced subsurface sequestration of supercritical CO_2 . *Int. J. Greenhouse Gas Control* **2009**, *3*, 90–99.
- (5) Mitchell, A. C.; Phillips, A. J.; Hamilton, M. A.; Gerlach, R.; Hollis, W. K.; Kaszuba, J. P.; Cunningham, A. B. Resilience of planktonic and biofilm cultures to supercritical CO_2 . *J. Supercrit. Fluids* **2009**, *47*, 318–325.
- (6) Gilfillan, S. M. V.; Lollar, B. S.; Holland, G.; Blagburn, D.; Stevens, S.; Schoell, M.; Cassidy, M.; Ding, Z. J.; Zhou, Z.; Lacrampe-Couloume, G.; Ballentine, C. J. Solubility trapping in formation water as dominant CO_2 sink in natural gas fields. *Nature* **2009**, *458* (7238), 614–618.
- (7) Stumm, W.; Morgan, J. J., *Aquatic Chemistry*, 3rd ed.; John Wiley & Sons: New York, 1996; p 1040.
- (8) Kaszuba, J. P.; Janecky, D. R., Geochemical impacts of sequestering carbon dioxide in brine aquifers. In *The Science and Technology of Carbon Sequestration*; McPherson, J. M.; Sundquist, E., Eds.; American Geophysical Union Monograph: Washington, DC, 2010; in press.
- (9) Gunter, W. D.; Perkins, E. H.; Hutcheon, I. Aquifer disposal of acid gases: modelling of water-rock reactions for trapping of acid wastes. *Appl. Geochem.* **2000**, *15* (8), 1085–1095.
- (10) Kaszuba, J. P.; Janecky, D. R.; Snow, M. G. Experimental evaluation of mixed fluid reactions between supercritical carbon dioxide and NaCl brine: Relevance to the integrity of a geologic carbon repository. *Chem. Geol.* **2005**, *217* (3–4), 277–293.
- (11) Schultze-Lam, S.; Fortin, D.; Davis, B. S.; Beveridge, T. J. Mineralization of bacterial surfaces. *Chem. Geol.* **1996**, *132* (1–4), 171–181.
- (12) Mitchell, A. C.; Ferris, F. G. The coprecipitation of Sr into calcite precipitates induced by bacterial ureolysis in artificial groundwater: Temperature and kinetic dependence. *Geochim. Cosmochim. Acta* **2005**, *69* (17), 4199–4210.
- (13) Van Lith, Y.; Warthmann, R.; Vasconcelos, C.; McKenzie, J. A. Microbial fossilization in carbonate sediments: A result of the bacterial surface involvement in dolomite precipitation. *Sedimentology* **2003**, *50*, 237–245.
- (14) Bell, P. E.; Mills, A. L.; Herman, J. S. Biogeochemical conditions favoring magnetite formation during anaerobic iron reduction. *Appl. Environ. Microbiol.* **1987**, *53* (11), 2610–2616.
- (15) Ferris, F. G.; Thompson, J. B.; Beveridge, T. J. Modern freshwater microbialites from Kelly lake, British Columbia, Canada. *Palaio* **1997**, *12* (3), 213–219.
- (16) Ferris, F. G.; Stehmeier, L. G.; Kantzas, A.; Mourits, F. M. Bacteriogenic mineral plugging. *J. Can. Pet. Technol.* **1996**, *35* (8), 56–61.
- (17) Mitchell, A. C.; Ferris, F. G. The Influence of *Bacillus pasteurii* on the nucleation and growth of calcium carbonate. *Geomicrobiol. J.* **2006**, *23* (3–4), 213–226.
- (18) Mitchell, A. C.; Ferris, F. G. Effect of strontium contaminants upon the size and solubility of calcite crystals precipitated by the bacterial hydrolysis of urea. *Environ. Sci. Technol.* **2006**, *40* (3), 1008–1014.
- (19) Fujita, Y.; Taylor, J. L.; Gresham, T. L.; Delwiche, M. E.; Colwell, F. S.; McLing, T.; Petzke, L.; Smith, R. W. Stimulation of microbial urea hydrolysis in groundwater to enhance calcite precipitation. *Environ. Sci. Technol.* **2008**, *42*, 3025–3032.

- (20) Busby, J. F.; Kimball, B. A.; Downey, J. S.; Peter, K. D. *Geochemistry of water in aquifers and confining units of the Northern Great Plains in parts of Montana, North Dakota, South Dakota, and Wyoming*, U.S.G.S. Professional Paper 1402-F; U.S. Geological Survey: Washington, DC, 1995; p 146.
- (21) Zhang, J.; Davis, T. A.; Matthews, M. A.; Drews, M. J.; LaBerge, M.; An, Y. H. Sterilization using high-pressure carbon dioxide. *J. Supercrit. Fluids* **2006**, *38* (3), 354–372.
- (22) Oldenburg, C. M.; Pruess, K.; Benson, S. M. Process Modeling of CO₂ injection into natural gas reservoirs for carbon sequestration and enhanced gas recovery. *Energy Fuels* **2001**, *15* (2), 293–298.
- (23) Parkhurst, D.; Appelo, C. *User's Guide to PHREEQC (Version 2)—A Computer Program for Speciation, Batch-Reaction, One-Dimensional Transport, And Inverse Geochemical Calculations*, Water-Resources Investigations Report 99-4259; U.S. Geological Survey: Washington, DC, 1999.
- (24) Allison, J. D.; Brown, D. S.; Novo-Gradac, K. J. *MINTEQA2/PRODEFA2—A Geochemical Assessment Model for Environmental Systems: Version 3.0 User's Manual*; U.S. Environmental Protection Agency: Washington, DC, 1990.
- (25) Stokes, R. Thermodynamics of aqueous urea solutions. *Aust. J. Chem.* **1967**, *20* (10), 2087–2100.
- (26) Fujita, Y.; Ferris, F. G.; Lawson, R. D.; Colwell, F. S.; Smith, R. W. Calcium carbonate precipitation by ureolytic subsurface bacteria. *Geomicrobiol. J.* **2000**, *17* (1), 305–318.
- (27) Hammes, F.; Boon, N.; de Villiers, J.; Verstraete, W.; Siciliano, S. D. Strain-specific ureolytic microbial calcium carbonate precipitation. *Appl. Environ. Microbiol.* **2003**, *69* (8), 4901–4909.
- (28) Warren, L. A.; Maurice, P. A.; Parmar, N.; Ferris, F. G. Microbially mediated calcium carbonate precipitation: Implications for interpreting calcite precipitation and for solid-phase capture of inorganic contaminants. *Geomicrobiol. J.* **2001**, *18* (1), 93–115.
- (29) Hoefs, J., *Stable Isotope Geochemistry*; Springer: Berlin, 1973; p 140.
- (30) Lesniak, P. M.; Zawidzki, P. Determination of carbon fractionation factor between aqueous carbonate and CO₂(g) in two-direction isotope equilibration. *Chem. Geol.* **2006**, *231* (3), 203–213.
- (31) Zhang, J.; Quay, P. D.; Wilbur, D. O. Carbon isotope fractionation during gas-water exchange and dissolution of CO₂. *Geochim. Cosmochim. Acta* **1995**, *59* (1), 107–114.
- (32) Ferris, F. G.; Fyfe, W. S.; Beveridge, T. J. Metallic Ion Binding by *Bacillus subtilis*—Implications for the Fossilization of Microorganisms. *Geology* **1988**, *16* (2), 149–152.
- (33) Wierzbowski, H. Effects of pre-treatments and organic matter on oxygen and carbon isotope analyses of skeletal and inorganic calcium carbonate. *Int. J. Mass Spectrom.* **2007**, *268* (1), 16–29.
- (34) Nonner, J., *Introduction to Hydrogeology*; Taylor & Francis: London, 2002; p 258.
- (35) Knobel, L. L.; Bartholomay, R. C.; Cecil, L. D.; Tucker, B. J.; Wegner, S. J., *Chemical Constituents in the Dissolved and Suspended Fractions of Ground Water from Selected Sites, Idaho National Engineering Laboratory and Vicinity, Idaho, 1989*; U.S. Geological Survey: Washington, DC, 1992; pp 92–51.
- (36) Carpenter, A. B.; Trout, M. L.; Pickett, E. E. Preliminary report on origin and chemical evolution of lead-rich and zinc-rich oil field brines in central Mississippi. *Econ. Geol.* **1974**, *69* (8), 1191–1206.
- (37) Richter, B.; Kreidler, C., *Geochemical Techniques for Identifying Sources of Ground-Water Salinization*; CRC Press: New York, 1993; p 258.
- (38) Lloyd, A. B.; Sheaffe, M. J. Urease activity in soils. *Plant Soil* **1973**, *39* (1), 71–80.
- (39) Mitchell, A.; Phillips, A.; Kaszuba, J.; Hollis, W.; Cunningham, A.; Gerlach, R. Microbially enhanced carbonate mineralization and the geologic containment of CO₂. *Geochim. Cosmochim. Acta* **2008**, *72* (12), A636.
- (40) World Water Assessment Programme, *Water for People: Water for Life - The UN World Water Development Report (The United Nations World Water Development Report)*; Berghahn Books: Oxford, NY, 2003.
- (41) Rittstiegl, K.; Robra, K. H.; Somitsch, W. Aerobic treatment of a concentrated urea wastewater with simultaneous stripping of ammonia. *Appl. Microbiol. Biotechnol.* **2001**, *56* (5), 820–825.

ES903270W